arXiv:0911.0236v1 [math.AG] 2 Nov 2009

On several families of elliptic curves with arbitrary large Selmer groups

Fei Li and Derong Qiu *
(School of Mathematical Sciences,
Institute of Mathematics and Interdisciplinary Science,
Capital Normal University, Beijing 100048, P.R.China)

1 Introduction and Main Results

In this paper, we consider the elliptic curve

$$E = E_{\varepsilon} : y^2 = x(x + \varepsilon pD)(x + \varepsilon qD), \quad (\varepsilon = \pm 1) \tag{1.1}$$

where p and q are odd prime numbers with $q - p = 2$, and $D = D_1 \cdots D_n$ is a square-free integer with distinct primes $D_1, \cdots, D_n$. Moreover, $2 \nmid D$, $p \nmid D$ and $q \nmid D$. For each D_i, denote $\widehat{D_i} = D/D_i$ ($\widehat{D_1} = 1$ if $D = D_1$). We write $E = E_+$ if $\varepsilon = 1$ and $E = E_-$ if $\varepsilon = -1$.

Take the elliptic curve

$$E' = E'_{\varepsilon} : y^2 = x^3 - 2\varepsilon(p+q)Dx^2 + 4D^2x. \tag{1.2}$$

There is an isogeny φ of degree 2 between E and E' as follows:

$$\varphi : E \longrightarrow E', \quad (x, y) \longmapsto (y^2/x^2,\ y(pqD^2 - x^2)/x^2).$$

The kernel is $E[\varphi] = \{O,\ (0,0)\}$, and the dual isogeny of φ is

$$\widehat{\varphi} : E' \longrightarrow E, \quad (x, y) \longmapsto (y^2/4x^2,\ y(4D^2 - x^2)/8x^2)$$

---

* E-mail: derong@mail.cnu.edu.cn

with kernel $E'[\widehat{\varphi}] = \{O, \ (0,0)\}$ (see [S, p.74]).

In this paper, we calculate the $\varphi(\widehat{\varphi})-$Selmer groups $S^{(\varphi)}(E/\mathbb{Q})$ and $S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ of the above elliptic curves via descent theory (see [S, Chapter X]), in particular, we obtain that the Selmer groups of several families of such elliptic curves can be arbitrary large. There are many literature studying Selmer groups of elliptic curves (see e.g., [KS], [QZ], [Sch], [ST]). The main results in the present paper are stated as follows:

Part 1 Results for the case $\varepsilon = +1$

Theorem 1.1. Let $E = E_+$ be the elliptic curve in (1.1) with $\varepsilon = +1$. For each $i \in \{1, \cdots, n\}$, denote

$$\Pi_i^+(D) = \left(1 - \left(\frac{-1}{D_i}\right)\right) + \left(1 - \left(\frac{p}{D_i}\right)\right) + \left(1 - \left(\frac{q}{D_i}\right)\right) + \sum_{j=1, \ j\neq i}^{n} \left(1 - \left(\frac{D_j}{D_i}\right)\right),$$

where $(-)$ is the (Legendre) quadratic residue symbol. And define a function $\rho^+(D)$ by

$$\rho^+(D) = \sum_{i=1}^{n} \left[\frac{1}{1 + \Pi_i^+(D)}\right],$$

where [x] is the greatest integer $\leq x$. Then we have

(1) There exists a subset $T \subset \{D_1, \cdots, D_n\}$ with cardinal $\sharp T = \rho^+(D)$ such that $S^{(\varphi)}(E/\mathbb{Q}) \supset < T \text{mod}(\mathbb{Q}^{\star^2}) > \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^+(D)}$. In particular, $\dim_2 S^{(\varphi)}(E/\mathbb{Q}) \geq \rho^+(D)$.

(2) If $p \equiv 7(\text{mod}8)$ and $D_i \equiv 1, \ 7(\text{mod}8)$ $(1 \leq i \leq n)$, then $\dim_2 S^{(\varphi)}(E/\mathbb{Q}) \geq \rho^+(D) + 1$ and $S^{(\varphi)}(E/\mathbb{Q}) \supset < T \cup \{2\} \text{mod}(\mathbb{Q}^{\star^2}) > \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^+(D)+1}$ for some subset $T \subset \{D_1, \cdots, D_n\}$ with $\sharp T = \rho^+(D)$.

Corollary 1.2. For elliptic curve $E = E_+$ as in Theorem 1.1, we have $S^{(\varphi)}(E/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+1}$. Furthermore,

(A) If the following conditions hold:

(1) $D_i \equiv 1(\mathrm{mod} 4), \quad 1 \leq i \leq n;$ (2) $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1, \quad 1 \leq i \leq n;$

(3) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \leq i, \ j \leq n$ and $i \neq j$.

Then $S^{(\varphi)}(E/\mathbb{Q}) \supset (\mathbb{Z}/2\mathbb{Z})^n$, so $2^n \leq \sharp S^{(\varphi)}(E/\mathbb{Q}) \leq 2^{n+1}$.

(B) If the following conditions hold:

(1) $D_i \equiv 1(\mathrm{mod} 4), \quad 1 \leq i \leq n;$

(2) $D_k \equiv 5(\mathrm{mod} 8)$ for at least one $k \in \{1, \cdots, n\}$;

(3) $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1, \quad 1 \leq i \leq n;$

(4) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \leq i, \ j \leq n$ and $i \neq j$.

Then $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^n$, and $\sharp S^{(\varphi)}(E/\mathbb{Q}) = 2^n$.

(C) If the following conditions hold:

(1) $p \equiv 7(\mathrm{mod} 8);$ (2) $D_i \equiv 1(\mathrm{mod} 8), \quad 1 \leq i \leq n;$

(3) $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1, \quad 1 \leq i \leq n;$

(4) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \leq i, \ j \leq n$ and $i \neq j$.

Then $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$, and $\sharp S^{(\varphi)}(E/\mathbb{Q}) = 2^{n+1}$.

Theorem 1.3. Let $E' = E'_+$ be the elliptic curve in (1.2) with $\varepsilon = +1$. For each $i \in \{1, \cdots, n\}$, denote

$$\Pi_i^+(D)' = \left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right)\left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) + \sum_{j=1, \ j\neq i}^{n} \left(1 - \left(\frac{D_i}{D_j}\right)\right)\left(1 - \left(\frac{pqD_i}{D_j}\right)\right),$$

where $(-)$ is the (Legendre) quadratic residue symbol. Take a subset I of $Z(n) = \{1, \cdots, n\}$ as follows: $I = \{i \in Z(n) : \ D_i \equiv 1(\mathrm{mod} 8)\} \cup \{i \in Z(n) : \ (1 + p\widehat{D_i})(1 +$

$q\widehat{D_i}) \equiv 0(\mathrm{mod}16)\} \cup \{i \in Z(n):\ D_i \equiv 3(\mathrm{mod}8)$ and $p \equiv 1(\mathrm{mod}4)\} \cup \{i \in Z(n):$ $D_i \equiv 7(\mathrm{mod}8)$ and $p \equiv 3(\mathrm{mod}4)\}$, and define a function $\rho^+(D)'$ by

$$\rho^+(D)' = \sum_{i\in I} \left[\frac{1}{1+\Pi_i^+(D)'}\right],$$

where [x] is the greatest integer $\leq x$. Then there exists a subset $T \subset \{D_1, \cdots, D_n\}$ with cardinal $\sharp T = \rho^+(D)'$ such that $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \supset < T\mathrm{mod}(\mathbb{Q}^{\star^2}) > \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^+(D)'}$. In particular, $\dim_2 S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \geq \rho^+(D)'$.

Corollary 1.4. For elliptic curve $E' = E'_+$ as in Theorem 1.3, we have $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+3}$. Furthermore, if for each $i \in \{1, \cdots, n\}$, $\Pi_i^+(D)' = 0$ and one of the following conditions hold:

(1) $D_i \equiv 1(\mathrm{mod}8)$; (2) $(1+p\widehat{D_i})(1+q\widehat{D_i}) \equiv 0(\mathrm{mod}16)$;

(3) $\left\{\begin{array}{l} D_i \equiv 3(\mathrm{mod}8), \\ p \equiv 1(\mathrm{mod}4); \end{array}\right.$ (4) $\left\{\begin{array}{l} D_i \equiv 7(\mathrm{mod}8), \\ p \equiv 3(\mathrm{mod}4). \end{array}\right.$

Then $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \supset (\mathbb{Z}/2\mathbb{Z})^{n+2}$, so $2^{n+2} \leq \sharp S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \leq 2^{n+3}$.

Furthermore, if $\left\{\begin{array}{l} p \equiv 3(\mathrm{mod}4), \\ \alpha(-pq)' = 0 \end{array}\right.$ or $\left\{\begin{array}{l} D-p \equiv 0, 2(\mathrm{mod}8), \\ \alpha(-pq)' = 0, \end{array}\right.$

then $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = (\mathbb{Z}/2\mathbb{Z})^{n+3}$, so $\sharp S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = 2^{n+3}$,

where $\alpha(-pq)' = \sum_{i=1}^n \left(1 - \left(\frac{-1}{D_i}\right)\right)\left(1 - \left(\frac{-pq}{D_i}\right)\right)$.

For example, if the following conditions hold:

(1) $D_i \equiv 1(\mathrm{mod}8)$, $1 \leq i \leq n$; (2) $\left(\frac{D_i}{p}\right) + \left(\frac{D_i}{q}\right) = 0$, $1 \leq i \leq n$;

(3) $p \equiv 3(\mathrm{mod}4)$.

Then $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = (\mathbb{Z}/2\mathbb{Z})^{n+3}$, so $\sharp S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = 2^{n+3}$.

Theorem 1.5. For elliptic curves $E = E_+$ and $E' = E'_+$in (1.1) and (1.2) with $\varepsilon = +1$, we have

(A) If the following conditions hold:

(1) $D_i \equiv 1(\mathrm{mod} 4), \quad 1 \le i \le n$;

(2) $D_k \equiv 5(\mathrm{mod} 8)$ for at least one $k \in \{1, \cdots, n\}$;

(3) $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1, \quad 1 \le i \le n$;

(4) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \le i,\ j \le n$ and $i \ne j$;

(5) $p - D \equiv 0, 2(\mathrm{mod} 8)$.

Then $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^n$, $(\mathbb{Z}/2\mathbb{Z})^{n+2} \subset S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+3}$.

Furthermore, if $p \equiv 3(\mathrm{mod} 4)$ or $p - D \equiv 0(\mathrm{mod} 8)$, then

$S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^n, \quad S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+3}$, and

$\mathrm{rank}(E(\mathbb{Q})) + \dim_2 \mathrm{TS}(E/\mathbb{Q})[\varphi] + \dim_2 \mathrm{TS}(E'/\mathbb{Q})[\widehat{\varphi}] = 2n + 1$,

where $\mathrm{TS}(E/\mathbb{Q})$ and $\mathrm{TS}(E'/\mathbb{Q})$ are the Tate-Shafarevich groups of E and E' over $\mathbb{Q}$ respectively (See [S, Chapter X]).

(B) If the following conditions hold:

(1) $p \equiv 7(\mathrm{mod} 8)$; (2) $D_i \equiv 1(\mathrm{mod} 8), \quad 1 \le i \le n$;

(3) $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1, \quad 1 \le i \le n$;

(4) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \le i,\ j \le n$ and $i \ne j$.

Then $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}, \quad S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+3}$, and

$\mathrm{rank}(E(\mathbb{Q})) + \dim_2 \mathrm{TS}(E/\mathbb{Q})[\varphi] + \dim_2 \mathrm{TS}(E'/\mathbb{Q})[\widehat{\varphi}] = 2n + 2$.

Part 2 Results for the case $\varepsilon = -1$

Theorem 1.6. Let $E = E_-$ be the elliptic curve in (1.1) with $\varepsilon = -1$. For each $i \in \{1, \cdots, n\}$, denote

$$\Pi_i^-(D) = \left(1 - \left(\frac{p}{D_i}\right)\right) + \left(1 - \left(\frac{q}{D_i}\right)\right) + \sum_{j=1,\ j\ne i}^{n} \left(1 - \left(\frac{D_j}{D_i}\right)\right),$$

and define a function $\rho^-(D)$ by

$$\rho^-(D) = \sum_{i=1}^{n} \left[\frac{1}{1+\Pi_i^-(D)}\right],$$

where [x] is the greatest integer $\leq x$. Then we have

(1) There exists a subset $T \subset \{D_1, \cdots, D_n\}$ with cardinal $\sharp T = \rho^-(D)$ such that $S^{(\varphi)}(E/\mathbb{Q}) \supset < \{D_i^\star : \ D_i \in T\} \mathrm{mod}(\mathbb{Q}^{\star^2}) > \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^-(D)}$, where $D_i^\star \in \{D_i, \ -D_i\}$. In particular, $\dim_2 S^{(\varphi)}(E/\mathbb{Q}) \geq \rho^-(D)$.

(2) Furthermore, if one of the following conditions hold:

(a) $p \equiv 7(\mathrm{mod} 8)$ and $D_i \equiv 1, \ 7(\mathrm{mod} 8)$ $(1 \leq i \leq n)$;

(b) $p \equiv 1(\mathrm{mod} 8)$ and $D_i \equiv 1, \ 3(\mathrm{mod} 8)$ $(1 \leq i \leq n)$.

Then for T as in (1), we have $S^{(\varphi)}(E/\mathbb{Q}) \supset < 2^\star > \times < \{D_i^\star : \ D_i \in T\} \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^-(D)+1}$, where $2^\star \in \{2, \ -2\}$. In particular, $\dim_2 S^{(\varphi)}(E/\mathbb{Q}) \geq \rho^-(D) + 1$.

Corollary 1.7. For elliptic curve $E = E_-$ as in Theorem 1.6, we have $S^{(\varphi)}(E/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+1}$. Furthermore,

(A) If the following conditions hold:

(1) $\left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1, \quad 1 \leq i \leq n;$

(2) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \leq i, \ j \leq n$ and $i \neq j$.

Then $S^{(\varphi)}(E/\mathbb{Q}) \supset (\mathbb{Z}/2\mathbb{Z})^n$, so $2^n \leq \sharp S^{(\varphi)}(E/\mathbb{Q}) \leq 2^{n+1}$.

(B) If the following conditions hold:

(1) $\left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1, \quad 1 \leq i \leq n;$

(2) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \leq i, \ j \leq n$ and $i \neq j$;

(3) $\begin{cases} p \equiv 7(\mathrm{mod} 8) \\ D_i \equiv 1, 7(\mathrm{mod} 8) \ (1 \leq i \leq n), \end{cases}$ or $\begin{cases} p \equiv 1(\mathrm{mod} 8) \\ D_i \equiv 1, 3(\mathrm{mod} 8) \ (1 \leq i \leq n). \end{cases}$

Then $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$, and $\sharp S^{(\varphi)}(E/\mathbb{Q}) = 2^{n+1}$.

Theorem 1.8. Let $E' = E'_-$ be the elliptic curve in (1.2) with $\varepsilon = -1$. For each $i \in \{1, \cdots, n\}$, denote

$$\Pi_i^-(D)' = \left(1 - \left(\frac{p\widehat{D_i}}{D_i}\right)\right)\left(1 - \left(\frac{q\widehat{D_i}}{D_i}\right)\right) + \sum_{j=1,\ j\neq i}^{n} \left(1 - \left(\frac{D_i}{D_j}\right)\right)\left(1 - \left(\frac{pqD_i}{D_j}\right)\right),$$

where $(-)$ is the (Legendre) quadratic residue symbol. Take a subset I of $Z(n) = \{1, \cdots, n\}$ as follows: $I = \{i \in Z(n):\ D_i \equiv 1(\mathrm{mod} 8)\} \cup \{i \in Z(n):\ (1 - p\widehat{D_i})(1 - q\widehat{D_i}) \equiv 0(\mathrm{mod} 16)\} \cup \{i \in Z(n):\ D_i \equiv 3(\mathrm{mod} 8)$ and $p \equiv 1(\mathrm{mod} 4)\} \cup \{i \in Z(n):$ $D_i \equiv 7(\mathrm{mod} 8)$ and $p \equiv 3(\mathrm{mod} 4)\}$, and define a function $\rho^-(D)'$ by

$$\rho^-(D)' = \sum_{i \in I} \left[\frac{1}{1 + \Pi_i^-(D)'}\right],$$

where [x] is the greatest integer $\leq x$. Then there exists a subset $T \subset \{D_1, \cdots, D_n\}$ with cardinal $\sharp T = \rho^-(D)'$ such that $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \supset < T\mathrm{mod}(\mathbb{Q}^{\star^2}) > \cong (\mathbb{Z}/2\mathbb{Z})^{\rho^-(D)'}$. In particular, $\dim_2 S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \geq \rho^-(D)'$.

Corollary 1.9. For elliptic curve $E' = E'_-$ as in Theorem 1.8, we have $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \subset (\mathbb{Z}/2\mathbb{Z})^{n+2}$. Furthermore, if for each $i \in \{1, \cdots, n\}$, $\Pi_i^-(D)' = 0$ and one of the following conditions hold:

(1) $D_i \equiv 1(\mathrm{mod} 8)$; (2) $(1 - p\widehat{D_i})(1 - q\widehat{D_i}) \equiv 0(\mathrm{mod} 16)$;

(3) $\begin{cases} D_i \equiv 3(\mathrm{mod} 8), \\ p \equiv 1(\mathrm{mod} 4); \end{cases}$ (4) $\begin{cases} D_i \equiv 7(\mathrm{mod} 8), \\ p \equiv 3(\mathrm{mod} 4). \end{cases}$

Then $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+2}$, so $\sharp S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = 2^{n+2}$.

For example, if the following conditions hold:

(1) $D_i \equiv 1(\mathrm{mod} 8)$, $1 \leq i \leq n$; (2) $\left(\frac{D_i}{p}\right) + \left(\frac{D_i}{q}\right) = 0$, $1 \leq i \leq n$.

Then $S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = (\mathbb{Z}/2\mathbb{Z})^{n+2}$, so $\sharp S^{(\widehat{\varphi})}(E'/\mathbb{Q}) = 2^{n+2}$.

Theorem 1.10 For elliptic curves $E = E_-$ and $E' = E'_-$ in (1.1) and (1.2) with $\varepsilon = -1$, we have

(A) If the following conditions hold:

(1) $D_i \equiv 1(\mathrm{mod}4), \quad 1 \leq i \leq n;$

(2) $D_k \equiv 5(\mathrm{mod}8)$ for at least one $k \in \{1, \cdots, n\}$;

(3) $\left(\frac{D_i}{p}\right) = \left(\frac{D_i}{q}\right) = 1, \quad 1 \leq i \leq n;$

(4) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \leq i,\ j \leq n$ and $i \neq j$;

(5) $p - D \equiv 2, 4(\mathrm{mod}8)$.

Then $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^n$, $\quad S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+2}$, and

$\mathrm{rank}(E(\mathbb{Q})) + \dim_2 \mathrm{TS}(E/\mathbb{Q})[\varphi] + \dim_2 \mathrm{TS}(E'/\mathbb{Q})[\widehat{\varphi}] = 2n.$

(B) If the following conditions hold:

(1) $p \equiv 1, 7(\mathrm{mod}8);$ $\qquad$ (2) $D_i \equiv 1(\mathrm{mod}8), \quad 1 \leq i \leq n;$

(3) $\left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1, \quad 1 \leq i \leq n;$

(4) $\left(\frac{D_j}{D_i}\right) = 1, \quad 1 \leq i,\ j \leq n$ and $i \neq j$.

Then $S^{(\varphi)}(E/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$, $\quad S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong (\mathbb{Z}/2\mathbb{Z})^{n+2}$, and

$\mathrm{rank}(E(\mathbb{Q})) + \dim_2 \mathrm{TS}(E/\mathbb{Q})[\varphi] + \dim_2 \mathrm{TS}(E'/\mathbb{Q})[\widehat{\varphi}] = 2n + 1.$

Remark 1.11 For each pair (p, q) above, by Chinese Remainder Theorem (See [L]) and Dirichlet Theorem on primes in arithmetic progressions [see[IK]], it is easy to verify that there exist infinitely many D satisfying the conditions in every Corollaries above.

2 Proofs of Theorems

Let $M_{\mathbb{Q}}$ be the set of all places of $\mathbb{Q}$, including the infinite ∞. For each place p, denote by $\mathbb{Q}_p$ the completion of $\mathbb{Q}$ at p, and if p is finite, denote by v_p the corresponding normalized additive valuation, so $v_p(p) = 1$. Let $S = \{\infty, 2, p, q, D_1, \cdots, D_n\}$, and define a subgroup of $\mathbb{Q}^\star/\mathbb{Q}^{\star^2}$ as follows: $\mathbb{Q}(S,2) =< -1, 2, p, q, D_1, \cdots, D_n >\cong (\mathbb{Z}/2\mathbb{Z})^{n+4}$. For any subset $A \subset \mathbb{Q}^\star$, we write $< A >$ for the subgroup of $\mathbb{Q}^\star/\mathbb{Q}^{\star^2}$ generated by all the elements in A. For each $d \in \mathbb{Q}(S,2)$, define the curves

$C_d: \ dw^2 = d^2 - 2\varepsilon(p+q)Ddz^2 + 4D^2z^4$, and

$C'_d: \ dw^2 = d^2 + \varepsilon(p+q)Ddz^2 + pqD^2z^4$.

Then by the theory of descent via two-isogeny (see [S, chapter X]), the $\varphi-$Selmer group for the elliptic curve $E/\mathbb{Q}$ in (1.1) is

$S^{(\varphi)}(E/\mathbb{Q}) \cong \{d \in \mathbb{Q}(S,2): \ C_d(\mathbb{Q}_v) \neq \emptyset \text{ for all } v \in S\}$, and the $\widehat{\varphi}-$Selmer group for the elliptic curve $E'/\mathbb{Q}$ in (1.2) is

$S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \cong \{d \in \mathbb{Q}(S,2): \ C'_d(\mathbb{Q}_v) \neq \emptyset \text{ for all } v \in S\}$.

Proposition 2.1 We assume $\varepsilon = 1$ and the elliptic curve $E = E_+$ be as in (1.1).

(A) For $d \in \mathbb{Q}(S,2)$, if one of the following conditions holds:

(1) $d < 0$; (2) $p \mid d$; (3) $q \mid d$.

Then $d \notin S^{(\varphi)}(E/\mathbb{Q})$. Moreover, if $d > 0$, then $C_d(\mathbb{R}) \neq \emptyset$.

(B) For $d = 2 \in \mathbb{Q}(S,2)$, we have

(1) $C_2(\mathbb{Q}_2) \neq \emptyset \iff D(D - 2p - 2) \equiv 1 (\text{mod} 16)$.

(2) For each prime number $l \mid pqD$, $C_2(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{2}{l}\right) = 1$, i.e., $l \equiv 1, 7 (\text{mod} 8)$.

(C) For each D_i $(1 \leq i \leq n)$, we have

(1) $C_{D_i}(\mathbb{Q}_2) \neq \emptyset \iff D_i \equiv 1 (\text{mod} 4)$.

(2) For each prime number $l \mid pq\widehat{D_i}$, $C_{D_i}(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{D_i}{l}\right) = 1$.

(3) $C_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(\frac{p\widehat{D_i}}{D_i}\right) = \left(\frac{q\widehat{D_i}}{D_i}\right) = 1$.

Proof. Conclusion (A) follows directly by the valuation property.

(B) In this case, $C_2 : w^2 = 2 - 2(p+q)Dz^2 + 2D^2z^4$.

(1) If $C_2(\mathbb{Q}_2) \neq \emptyset$, then there exists $(z_0, w_0) \in \mathbb{Q}_2^2$ such that $w_0^2 = 2 - 2(p+q)Dz_0^2 + 2D^2z_0^4$, from which we have $v_2(w_0) \geq 1$ and $v_2(z_0) = 0$. Let $w_0 = 2w_1$ with $w_1 \in \mathbb{Q}_2$ and $v_2(w_1) \geq 0$. Then $2w_1^2 = 1 - (p+q)Dz_0^2 + D^2z_0^4$. Note that $v_2(Dz_0) = 0$, so by lemma in [Rob], we have $z_0^2 = 8s+1$ and $D^2z_0^4 = 8t+1$ for some $s, t \in \mathbb{Q}_2$ with $v_2(s) \geq 0$ and $v_2(t) \geq 0$. So $2w_1^2 = 2 - 2(p+1)D(8s+1) + 8t$. Thus $v_2(2w_1^2) = v_2(2) = 1$, so $v_2(w_1) = 0$, and we may assume that $(w_1)^2 = 8r+1$ for some $r \in \mathbb{Q}_2$ and $v_2(r) \geq 0$. So $2(8r+1) = 1 - 2(p+1)D(8s+1) + D^2(8s+1)^2$, from which we get $D^2 - 2(p+1)D \equiv 1(\mathrm{mod}16)$, i.e., $D(D-2p-2) \equiv 1(\mathrm{mod}16)$.

Conversely, if $D(D-2p-2) \equiv 1(\mathrm{mod}16)$, let $f(z,w) = w^2 - 2 + 2(p+q)Dz^2 - 2D^2z^4$, then $f'_w = 2w$, and $v_2(f(1,2)) \geq 5 > 2v_2(f'_w(1,2))$, so by Hensel's lemma (see [S],*p*.322), we have $C_2(\mathbb{Q}_2) \neq \emptyset$.

(2) For $l = p$, if $C_2(\mathbb{Q}_p) \neq \emptyset$, then $\exists (z_0, w_0) \in \mathbb{Q}_p^2$ s.t. $w_0^2 = 2 - 2(p+q)Dz_0^2 + 2D^2z_0^4$. If $v_p(z_0) > 0$, then $v_p(w_0^2 - 2) \geq 1$, so $\left(\frac{2}{p}\right) = 1$. If $v_p(z_0) = 0$ then we can obtain that $v_p(Dz_0^2 - 1) = 0$. Let $w_1 = w_0(Dz_0^2 - 1)^{-1} \in \mathbb{Q}_p^*$, then $v_p(w_1^2 - 2) \geq 1$, so $\left(\frac{2}{p}\right) = 1$. If $v_p(z_0) < 0$ then $v_p(z_0^{-1}) > 0$, so by the equality $(\frac{w_0}{z_0^2})^2 = 2(z_0^{-1})^4 - 4(p+1)Dz_0^{-2} + 2D^2$ we get $v_p(w_1) \geq 0$ and $v_p(w_1^2 - 2D^2) \geq 1$ with $w_1 = \frac{w_0}{z_0^2}$. So $\left(\frac{2D^2}{p}\right) = 1$, and then $\left(\frac{2}{p}\right) = 1$. To sum up, we obtain $\left(\frac{2}{p}\right) = 1$.

Conversely, if $\left(\frac{2}{p}\right) = 1$, then $2 \equiv a^2(\mathrm{mod}p)$ for some $a \in \mathbb{Z}$ with $p \nmid a$. Then for the above f(z, w), we have $v_p(f(0,a)) \geq 1 > 0 = 2v_p(f'_w(0,a))$. So by Hensel's lemma , we have $C_2(\mathbb{Q}_p) \neq \emptyset$. The case $l = q, D_i$ $(1 \leq i \leq n)$ are similar. This proves (B).

(C) In this case, $C_{D_i} : w^2 = D_i - 2(p+q)Dz^2 + 4D\widehat{D_i}z^4$.

(1) If $C_{D_i}(\mathbb{Q}_2) \neq \emptyset$, then $\exists (z_0, w_0) \in \mathbb{Q}_2^2$ s. t. $w_0^2 = D_i - 2(p+q)Dz_0^2 + 4D\widehat{D_i}z_0^4$.

If $v_2(z_0) \geq 0$, then $v_2(w_0^2) = v_2(D_i)$, $v_2(w_0) = 0$ and $v_2(w_0^2 - D_i) \geq 2$. Since $v_2(w_0^2 - D_i) \geq 3$ (see lemma in [Rob]), we have $v_2(D_i - 1) = v_2((w_0^2 - 1) + (D_i - w_0^2)) \geq 2$. So $D_i \equiv 1 (\mathrm{mod} 4)$.

If $v_2(z_0) < 0$, then $v_2(z_0^{-1}) \geq 1$, so by $(\frac{w_0}{2z_0^2})^2 = D_i(2z_0^2)^{-2} - (p+1)Dz_0^{-2} + D\widehat{D_i}$, we get $v_2((\frac{w_0}{2z_0^2})^2) = v_2(D\widehat{D_i}) = 0$, so $v_2((\frac{w_0}{2z_0^2}) = 0$ and $v_2(D_i\widehat{D_i}^2 - (\frac{w_0}{2z_0^2})^2) \geq 2$, hence $v_2(D_i - (\frac{w_0}{2z_0^2\widehat{D_i}})^2) \geq 2$. Since $v_2((\frac{w_0}{2z_0^2\widehat{D_i}})^2 - 1) \geq 3$, we get $v_2(D_i - 1) = v_2((D_i - (\frac{w_0}{2z_0^2\widehat{D_i}})^2) + ((\frac{w_0}{2z_0^2\widehat{D_i}})^2 - 1)) \geq 2$. So $D_i \equiv 1 (\mathrm{mod} 4)$.

Conversely, if $D_i \equiv 1 (\mathrm{mod} 4)$, then for the polynomial $f(z, w) = w^2 - D_i + 2(p+q)Dz^2 - 4D\widehat{D_i}z^4$, we have $f'_w = 2w$. So if $D_i \equiv 1 (\mathrm{mod} 8)$, then $v_2(f(0,1)) \geq 3 > 2 = 2v_2(f'_w(0,1))$; If $D_i \equiv 5 (\mathrm{mod} 8)$, then $v_2(f(1,1)) \geq 3 > 2 = 2v_2(f'_w(1,1))$. So by Hensel's lemma, we have $C_{D_i}(\mathbb{Q}_2) \neq \emptyset$.

(2) For $l = p$, if $C_{D_i}(\mathbb{Q}_p) \neq \emptyset$, then $\exists (z_0, w_0) \in \mathbb{Q}_p^2$ s. t. $w_0^2 = D_i - 2(p+q)Dz_0^2 + 4D\widehat{D_i}z_0^4 = -4pDz_0^2 + D_i(2\widehat{D_i}z_0^2 - 1)^2$.

If $v_p(z_0) > 0$, then $v_p(w_0^2) = v_p(D_i) = 0$, so $v_p(w_0) = 0$ and $v_p(w_0^2 - D_i) \geq 1$, so $\left(\frac{D_i}{p}\right) = 1$.

If $v_p(z_0) = 0$, then $v_p(w_0) \geq 0$. If $v_p(2\widehat{D_i}z_0^2 - 1) \geq 1$, then $v_p((2\widehat{D_i}z_0^2 - 1))^2 \geq 2 > 1 = v_p(-4pDz_0^2)$, so $v_p(w_0^2) = v_p(-4pDz_0^2)$, i.e., $2v_p(w_0^2) = 1$, which is impossible ! So $v_p(2\widehat{D_i}z_0^2 - 1) = 0$, and then we have $v_p(D_i - (\frac{w_0}{2\widehat{D_i}z_0^2 - 1})^2) \geq 1$, which shows that $\left(\frac{D_i}{p}\right) = 1$.

If $v_p(z_0) < 0$, then $v_p(z_0^{-1}) \geq 1$. By the above equation of C_{D_i} we have $(\frac{w_0}{2\widehat{D_i}z_0^2})^2 = D_i(2\widehat{D_i}z_0^2)^{-2} - \frac{(p+1)D}{(\widehat{D_i}z_0^2)^2} + D_i$. So $v_p((\frac{w_0}{2\widehat{D_i}z_0^2})^2) = v_p(D_i) = 0$, hence $v_p(D_i - \frac{w_0}{2\widehat{D_i}z_0^2}) \geq 1$, which shows $\left(\frac{D_i}{p}\right) = 1$.

Conversely, if $\left(\frac{D_i}{p}\right) = 1$, then $D_i \equiv a^2 (\mathrm{mod} p)$ for some $a \in \mathbb{Z}$ with $v_p(a) = 0$. Then $v_p(f(0,a)) \geq 1 > 0 = 2v_p(f'_w(0,a))$. By Hensel's lemma, we get $C_{D_i}(\mathbb{Q}_p) \neq \emptyset$. The other cases follow the same line. We omit the details. The proof of Proposition 2.1 is completed. $\square$

Corollary 2.2 We assume $\varepsilon = 1$ and the elliptic curve $E = E_+$ be as in (1.1).

(A) $2 \in S^{(\varphi)}(E/\mathbb{Q}) \iff p \equiv 7 (\mathrm{mod} 8)$ and $D_i \equiv 1, 7 (\mathrm{mod} 8)$ $(1 \leq i \leq n)$.

(B) For each D_i $(1 \leq i \leq n)$, we have $D_i \in S^{(\varphi)}(E/\mathbb{Q})$ if and only if $D_i \equiv 1 (\mathrm{mod} 4)$ and $\left(\frac{D_j}{D_i}\right) = \left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1$ $(1 \leq j \leq n,$ and $j \neq i)$.

Proof. The results follow easily from Proposition 2.1. $\square$

Proposition 2.3 We assume $\varepsilon = 1$ and the elliptic curve $E' = E'_+$ be as in (1.2).

(A) (1) For any $d \in \mathbb{Q}(S, 2)$, $C'_d(\mathbb{R}) \neq \emptyset$. If $2 \mid d$, then $d \notin S^{(\widehat{\varphi})}(E'/\mathbb{Q})$.

(2) $\{1, pq, -pD, -qD\} \subset S^{(\widehat{\varphi})}(E'/\mathbb{Q})$.

(B) For each D_i $(1 \leq i \leq n)$, we have

(1) $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$ if and only if one of the following conditions holds:

(a) $D_i \equiv 1 (\mathrm{mod} 8)$; (b) $(1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0 (\mathrm{mod} 16)$;

(c) $D_i \equiv 3 (\mathrm{mod} 8)$ and $p \equiv 1 (\mathrm{mod} 4)$; (d) $D_i \equiv 7 (\mathrm{mod} 8)$ and $p \equiv 3 (\mathrm{mod} 4)$.

(2) $C'_{D_i}(\mathbb{Q}_p) \neq \emptyset$ and $C'_{D_i}(\mathbb{Q}_q) \neq \emptyset$.

(3) For each $j \neq i$, $C'_{D_i}(\mathbb{Q}_{D_j}) \neq \emptyset \iff \left(1 - \left(\frac{D_i}{D_j}\right)\right)\left(1 - \left(\frac{pqD_i}{D_j}\right)\right) = 0$.

(4) $C'_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right)\left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) = 0$.

(C) For $d = -pq \in \mathbb{Q}(S, 2)$, we have

(1) $C'_{-pq}(\mathbb{Q}_2) \neq \emptyset \iff p \equiv 3 (\mathrm{mod} 4)$ or $D - p \equiv 0, 2 (\mathrm{mod} 8)$.

(2) $C'_{-pq}(\mathbb{Q}_p) \neq \emptyset$ and $C'_{-pq}(\mathbb{Q}_q) \neq \emptyset$.

(3) For each D_i $(1 \leq i \leq n)$, we have

$C'_{-pq}(\mathbb{Q}_{D_i}) \neq \emptyset \Longleftrightarrow \left(1 - \left(\frac{-1}{D_i}\right)\right)\left(1 - \left(\frac{-pq}{D_i}\right)\right) = 0.$

Proof. (A) (1) Obvious.

(2) For $C'_d : dw^2 = (d + pDz^2)(d + qDz^2)$, if $d = -pD$, then $(1, 0) \in C'_d(\mathbb{Q})$; and if $d = -qD$, then $(1, 0) \in C'_d(\mathbb{Q})$, so the group $< -pD, -qD >= \{1, pq, -pD, -qD\} \subseteq E(\mathbb{Q})/\widehat{\varphi}(E'(\mathbb{Q})) \subset S^{(\widehat{\varphi})}(E'/\mathbb{Q})$.

(B) C'_{D_i} : $w^2 = D_i + (p + q)Dz^2 + pqD\widehat{D_i}z^4 = D_i(1 + p\widehat{D_i}z^2)(1 + q\widehat{D_i}z^2)$.

(1) If $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$, then $\exists (z_0, w_0) \in \mathbb{Q}^2_{D_i}$ s. t. $w_0^2 = D_i + (p + q)Dz_0^2 + pqD\widehat{D_i}z_0^4$.

If $v_2(z_0) > 0$, then $v_2(w_0) = 0$. By Lemma 4 in [C, p.11], we get $v_2(D_i - 1) \geq 3$ and then $D_i \equiv 1 (\text{mod} 8)$.

If $v_2(z_0) \leq 0$, then by the equality $(\frac{w_0}{z_0^2})^2 = D_i(z_0^{-2} + p\widehat{D_i})(z_0^{-2} + q\widehat{D_i})$, we have $v_2(\frac{w_0}{z_0^2}) \geq 0$. Denote $W = \frac{w_0}{z_0^2}$, $Z = z_0^{-1}$, then $v_2(Z) \geq 0$, $v_2(W) \geq 0$ and $W^2 = D_i(Z^2 + p\widehat{D_i})(Z^2 + q\widehat{D_i}) = D_i(V^2 - \widehat{D_i}^2)$ with $V = Z^2 + (p + 1)\widehat{D_i} \in \mathbb{Q}_2$. Obviously, $v_2(V) \geq 0$.

i) If $v_2(V) = 0$, then $v_2(V^2 - 1) \geq 3$ and $v_2(Z) = 0$. So $v_2(V^2 - \widehat{D_i}^2) = v_2((V^2 - 1) - (\widehat{D_i}^2 - 1)) \geq 3$. Thus $v_2(W^2) = v_2(V^2 - \widehat{D_i}^2) \geq 3$, so $v_2(W^2) \geq 2$, and then $v_2((1 + p\widehat{D_i})(1 + q\widehat{D_i})) \geq 4$, i.e., $(1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0 (\text{mod} 16)$.

ii) If $v_2(V) \geq 1$, then $v_2(W) = 0$.

If $v_2(V) = 1$, then $v_2(V^2 - 4) \geq 5$. So $v_2(W^2 - 3D_i) = v_2((V^2 - 4) - (\widehat{D_i}^2 - 1)) \geq 3$. Thus $v_2(1 - 3D_i) \geq 3$, hence $D_i \equiv 3 (\text{mod} 8)$. Since $V = Z^2 + (p + 1)\widehat{D_i}$, we get $v_2(p + 1) = v_2(V - Z^2) = 1$, so $p \equiv 1 (\text{mod} 4)$.

If $v_2(V) \geq 2$, then $v_2(V^2) \geq 4$, so $W^2 + D_i = D_i(V^2 + 1 - \widehat{D_i}^2)$, hence $v_2(W^2 + D_i) \geq 3$, and then $v_2(1+D_i) = v_2(W^2+D_i+1-W^2) \geq 3$, so $D_i \equiv 7(\mathrm{mod}8)$. Since $V = Z^2+(p+1)\widehat{D_i}$, we get $v_2(p+1) = v_2(V-Z^2) \geq 2$, which shows $p \equiv 3(\mathrm{mod}4)$. To sum up, we have proved that if $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$, then one of the following conditions holds:

(a) $D_i \equiv 1(\mathrm{mod}8)$; (b) $(1+p\widehat{D_i})(1+q\widehat{D_i}) \equiv 0(\mathrm{mod}16)$;

(c) $D_i \equiv 3(\mathrm{mod}8)$ and $p \equiv 1(\mathrm{mod}4)$; (d) $D_i \equiv 7(\mathrm{mod}8)$ and $p \equiv 3(\mathrm{mod}4)$.

Conversely, we assume one of the above conditions holds. Let $f(z,\ w) = D_i+(p+q)Dz^2+pqD\widehat{D_i}z^4-w^2$, then $f'_w(z,w) = -2w$.

(a) If $D_i \equiv 1(\mathrm{mod}8)$, then $v_2(f(0,1)) \geq 3 > 2 = 2v_2(f'_w(0,1))$. So by Hensel's lemma, we get $C'_{D_i}(\mathbb{Q}_2) \neq \emptyset$.

(b) If $(1+p\widehat{D_i})(1+q\widehat{D_i}) \equiv 0(\mathrm{mod}16)$, then $v_2(1+p\widehat{D_i}) \geq 3$ or $v_2(1+q\widehat{D_i}) \geq 3$. Let $g(z) = 1+l\widehat{D_i}z^2$, where $l = p$ or q according to the above first or second inequalities. Then $g'(z) = 2l\widehat{D_i}z$. So $v_2(g(1)) \geq 3 > 2 = 2v_2(g'(1))$, and by Hensel's lemma, $g(z_0) = 0$ for some $z_0 \in \mathbb{Q}_2$. Therefore $(z_0, 0) \in C'_{D_i}(\mathbb{Q}_2)$.

(c) If $D_i \equiv 3(\mathrm{mod}8)$ and $p \equiv 1(\mathrm{mod}4)$, let $g(w) = w^2-D_i(2^2+p\widehat{D_i})(2^2+q\widehat{D_i})$, then $g'(w) = 2w$, and $v_2(g(1)) > 2 = 2v_2(g'(1))$. So by Hensel's lemma, $g(w_0) = 0$ for some $w_0 \in \mathbb{Q}_2$. Thus $(\frac{1}{2},\ \frac{w_0}{4}) \in C'_{D_i}(\mathbb{Q}_2)$.

(d) similar to (c) above. This proves (1).

(2) Let $f(z) = D_i+(p+q)Dz^2+pqD\widehat{D_i}z^4$, then $f(z) \equiv D_i(1+2\widehat{D_i}z^2)(\mathrm{mod}p)$, also $f(z) \equiv D_i(1-2\widehat{D_i}z^2)(\mathrm{mod}q)$. Take $\overline{g} = 1,\ \overline{h} = \overline{f}(\mathrm{mod}p)$ (if D_i is a square mod$p$, then take $\overline{g} = D_i(\mathrm{mod}p)$), then $\deg\overline{g} = 0$ and it is easy to verify that $\overline{h}$ is square free. So by lemma 14 in [MSS], $w^2 = f(z)$ has solutions in $\mathbb{Q}_p$, i.e., $C'_{D_i}(\mathbb{Q}_p) \neq \emptyset$. Similarly, $C'_{D_i}(\mathbb{Q}_q) \neq \emptyset$. This proves (2).

(3) If $C'_{D_i}(\mathbb{Q}_{D_j}) \neq \emptyset$ $(j \neq i)$, then $\exists (z_0, w_0) \in \mathbb{Q}^2_{D_j}$ s. t. $w_0^2 = D_i(1 + p\widehat{D_i}z_0^2)(1 + q\widehat{D_i}z_0^2)$.

i) If $v_{D_j}(z_0) \geq 0$, then $v_{D_j}(w_0) = 0$ and $D_i \equiv w_0^2 (\mathrm{mod} D_j)$, so $\left(\frac{D_i}{D_j}\right) = 1$.

ii) If $v_{D_j}(z_0) < 0$, then $v_{D_j}(z_0^{-1}) \geq 1$ and $v_{D_j}(w_0) = 1 + 2v_{D_j}(z_0) < 0$. Now $(\frac{w_0}{z_0^2})^2 = D_i z_0^{-4} + (p+q)Dz_0^{-2} + pqD_i\widehat{D_i}^2$. By the above discussion, we get $pqD_i \equiv (\frac{w_0}{z_0^2\widehat{D_i}})^2 (\mathrm{mod} D_j)$ with $v_{D_j}(\frac{w_0}{z_0^2\widehat{D_i}})$, so $\left(\frac{pqD_i}{D_j}\right) = 1$. To sum up, we get $\left(1 - \left(\frac{D_i}{D_j}\right)\right)\left(1 - \left(\frac{pqD_i}{D_j}\right)\right) = 0$.

Conversely, if $\left(1 - \left(\frac{D_i}{D_j}\right)\right)\left(1 - \left(\frac{pqD_i}{D_j}\right)\right) = 0$, then $D_i \equiv a^2 (\mathrm{mod} D_j)$ or $pqD_i \equiv b^2 (\mathrm{mod} D_j)$ for some $a, b \in \mathbb{Z}$ and $D_j \nmid ab$.

For the former, let $f(z, w) = w^2 - D_i - (p+q)Dz^2 + pqD_i\widehat{D_i}^2 z^4$, then $f'_w(z, w) = 2w$ and $v_{D_j}(f(0, a)) > 0 = 2v_{D_j}(f'_w(0, a))$. So by Hensel's lemma, $C'_{D_i}(\mathbb{Q}_{D_j}) \neq \emptyset$.

For the later, let $g(z, w) = w^2 - D_i\widehat{D_i}^2 - (p+q)\widehat{D_i}^2 Dz^2 - pqD_i\widehat{D_i}^4 z^4$, then $g'_w(z, w) = 2w$ and $v_{D_j}(g(\widehat{D_i}^{-1}, b)) > 0 = 2v_{D_j}(g'_w(\widehat{D_i}^{-1}, b))$. So by Hensel's lemma, $\exists (z_0, w_0) \in \mathbb{Q}^2_{D_j}$ s. t. $g(z_0, w_0) = 0$, and so $(z_0, \frac{w_0}{\widehat{D_i}}) \in C'_{D_i}(\mathbb{Q}_{D_j})$. This proves (3).

(4) If $C'_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset$, then $\exists (z_0, w_0) \in \mathbb{Q}^2_{D_i}$ s. t. $w_0^2 = D_i(1 + p\widehat{D_i}z_0^2)(1 + q\widehat{D_i}z_0^2)$. It is easy to see that $v_{D_i}(z_0) = 0$ and $v_{D_i}(w_0) > 0$. Set $w_0 = D_i w_1$ with $w_1 \in \mathbb{Q}_{D_i}$ and $v_{D_i}(w_1) \geq 0$. Then $D_i w_1^2 = (1 + p\widehat{D_i}z_0^2)(1 + q\widehat{D_i}z_0^2)$, so $(1 + p\widehat{D_i}z_0^2)(1 + q\widehat{D_i}z_0^2) \equiv 0 (\mathrm{mod} D_i)$. Thus $\left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right)\left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) = 0$.

Conversely, if $\left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right)\left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) = 0$, then $\exists a \in \mathbb{Z}$ such that $-l\widehat{D_i} \equiv a^2 (\mathrm{mod} D_i)$ for $l = p$ or q. Obviously, $D_i \nmid a$. Let $g(z) = 1 + l\widehat{D_i}z^2$, then $g'(z) = 2l\widehat{D_i}z$. So $v_{D_i}(g(a^{-1})) > 0 = 2v_{D_i}(g'(a^{-1}))$, by Hensel's lemma, $g(z_0) = 0$ for some $z_0 \in \mathbb{Q}_{D_i}$. Thus $(z_0, 0) \in C'_{D_i}(\mathbb{Q}_{D_i})$, so $C'_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset$. This proves (4), and (B) is proved.

(C) (1) If $C'_{-pq}(\mathbb{Q}_2) \neq \emptyset$, then $\exists (z_0, w_0) \in \mathbb{Q}^2_2$ s. t. $w_0^2 = -pq + (p+q)Dz_0^2 - D^2 z_0^4$.

i) If $v_2(z_0) < 0$, then $v_2(z_0^{-1}) \geq 1$, so by the equality $r_0^2 = -pqz_0^{-4} + (p+q)Dz_0^{-2} - D^2$ with $r_0 = \frac{w_0}{z_0^2}$, we get $v_2(r_0) = 0,\ v_2(r_0^2 + D^2) \geq 4$. Thus $v_2(r_0^2 + D^2) = v_2((r_0^2 - D^2) + 2D^2) = 1$, a contradiction! So we must have $v_2(z_0) \geq 0$.

ii) If $v_2(z_0) > 0$, then $v_2(w_0) = 0,\ v_2(w_0^2 + pq) \geq 4$. So we get $v_2(1 + pq) = v_2(1 - w_0^2 + w_0^2 + pq) \geq 3$, which shows $p \equiv 3(\mathrm{mod} 4)$.

iii) If $v_2(z_0) = 0$, then $v_2(z_0^2 - 1) \geq 3,\ v_2(z_0^4 - 1) \geq 4$. Since $w_0^2 = -(D-p)(D-q) + (p+q)D(z_0^2 - 1) - D^2(z_0^4 - 1)$, we have $v_2(w_0^2 + (D-p)(D-q)) \geq 4$. But $(D-p)(D-q) \equiv 0(\mathrm{mod} 8)$, so $v_2(w_0) \geq 2$. Thus $v_2((D-p)(D-q)) \geq 4$, which shows $D - p \equiv 0,\ 2(\mathrm{mod} 8)$.

Conversely, if $p \equiv 3(\mathrm{mod} 4)$, we get $-pq \equiv 1(\mathrm{mod} 8)$. So $-pq = a^2$ for some $a \in \mathbb{Q}_2$. Hence $(0, a) \in C'_{-pq}(\mathbb{Q}_2)$.

If $D - p \equiv 0,\ 2(\mathrm{mod} 8)$, then $(D - p - 1)^2 \equiv 1(\mathrm{mod} 16)$. Let $f(z, w) = -[(p+1)z^2 - D]^2 + z^4 - w^2$, then $f'_z(z, w) = -4(p+1)[(p+1)z^2 - D]z + 4z^3$.

If $v_2((D-p-1)^2 - 1) > 4$, then $v_2(f(1, 0)) > 4 = 2v_2(f'_z(1, 0))$;

If $v_2((D-p-1)^2 - 1) = 4$, then $v_2(f(3, 0)) \geq 5 > 4 = 2v_2(f'_z(3, 0))$. So by Hensel's lemma, $\exists (z_0, w_0) \in \mathbb{Q}_2^2$ s. t. $f(z_0, w_0) = 0$. It is easy to prove $z_0 \neq 0$. Hence $(z_0^{-1}, w_0 z_0^{-1}) \in C'_{-pq}(\mathbb{Q}_2)$.

(2) is similar to that of (B)(2).

(3) If $C'_{-pq}(\mathbb{Q}_{D_i}) \neq \emptyset$, then $\exists (z_0, w_0) \in \mathbb{Q}_{D_i}^2$ s. t. $w_0^2 = -pq + (p+q)Dz_0^2 - D^2 z_0^4$.

i) If $v_{D_i}(z_0) \geq 0$, then we get $\left(\frac{-pq}{D_i}\right) = 1$.

ii) If $v_{D_i}(z_0) < 0$, then $v_{D_i}(z_0^{-1}) \geq 1$. Since $(\frac{w_0}{Dz_0^2})^2 = -D^{-2}pqz_0^{-4} + (p+q)D^{-1}z_0^{-2} - 1$, we have $\left(\frac{-1}{D_i}\right) = 1$. To sum up, we get $\left(1 - \left(\frac{-1}{D_i}\right)\right)\left(1 - \left(\frac{-pq}{D_i}\right)\right) = 0$.

Conversely, if $\left(1 - \left(\frac{-1}{D_i}\right)\right)\left(1 - \left(\frac{-pq}{D_i}\right)\right) = 0$, then $\exists a, b \in \mathbb{Z}$ s. t. $-pq \equiv a^2(\mathrm{mod} D_i)$

or $-1 \equiv b^2 (\mathrm{mod} D_i)$ with $D_i \nmid ab$. For the former, let $f(z,w) = w^2 + pq - (p+q)Dz^2 + D^2 z^4$, then $f'_w(z,w) = 2w$ and $v_{D_i}(f(0,a)) > 0 = 2v_{D_i}(f'_w(0,a))$. So by Hensel's lemma, we get $C'_{-pq}(\mathbb{Q}_{D_i}) \neq \emptyset$.

For the later, let $f(z,w) = w^2 + D^2 pq - (p+q)D^3 z^2 + D^4 z^4$, then $f'_w(z,w) = 2w$ and $v_{D_i}(f(D^{-1},b)) > 0 = 2v_{D_i}(f'_w(D^{-1},b))$. So by Hensel's lemma, $\exists (z_0, w_0) \in \mathbb{Q}^2_{D_i}$ s. t. $f(z_0,w_0) = 0$. Thus $\exists (z_0, w_0 D^{-1}) \in C'_{-pq}(\mathbb{Q}_{D_i})$. This proves (C), and the proof of proposition 2.3 is completed. $\square$

Corollary 2.4 We assume $\varepsilon = 1$ and the elliptic curve $E' = E'_+$ be as in (1.2).

(A) Denote $\alpha(-pq)' = \sum_{i=1}^{n} \left(1 - \left(\frac{-1}{D_i}\right)\right)\left(1 - \left(\frac{-pq}{D_i}\right)\right)$. Then

$$-pq \in S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \iff \begin{cases} p \equiv 3 (\mathrm{mod} 4), \\ \alpha(-pq)' = 0; \end{cases} \text{or} \begin{cases} D - p \equiv 0, 2 (\mathrm{mod} 8), \\ \alpha(-pq)' = 0. \end{cases}$$

(B) Denote $\beta(-D)' = \sum_{i=1}^{n} \left(1 - \left(\frac{p}{D_i}\right)\right)\left(1 - \left(\frac{q}{D_i}\right)\right)$. Then $-D \in S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ if and only if one of the following conditions holds:

$$(1) \begin{cases} D \equiv 7 (\mathrm{mod} 8), \\ \beta(-D)' = 0; \end{cases} \qquad (2) \begin{cases} p \equiv 1, 7 (\mathrm{mod} 8), \\ \beta(-D)' = 0; \end{cases}$$

$$(3) \begin{cases} D \equiv 1 (\mathrm{mod} 8), \\ p \equiv 3 (\mathrm{mod} 8), \\ \beta(-D)' = 0; \end{cases} \qquad (4) \begin{cases} D \equiv 5 (\mathrm{mod} 8), \\ p \equiv 5 (\mathrm{mod} 8), \\ \beta(-D)' = 0. \end{cases}$$

(C) For each D_i $(1 \leq i \leq n)$, denote

$$\Pi_i^+(D)' = \left(1 - \left(\frac{-p\widehat{D_i}}{D_i}\right)\right)\left(1 - \left(\frac{-q\widehat{D_i}}{D_i}\right)\right) + \sum_{j=1,\ j\neq i}^{n} \left(1 - \left(\frac{D_i}{D_j}\right)\right)\left(1 - \left(\frac{pqD_i}{D_j}\right)\right).$$

Then $D_i \in S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ if and only if one of the following conditions holds:

$$(1) \begin{cases} D_i \equiv 1 (\mathrm{mod} 8), \\ \Pi_i^+(D)' = 0; \end{cases} \qquad (2) \begin{cases} (1 + p\widehat{D_i})(1 + q\widehat{D_i}) \equiv 0 (\mathrm{mod} 16), \\ \Pi_i^+(D)' = 0; \end{cases}$$

$$(3) \begin{cases} D_i \equiv 3 (\mathrm{mod} 8), \\ p \equiv 1 (\mathrm{mod} 4), \\ \Pi_i^+(D)' = 0; \end{cases} \qquad (4) \begin{cases} D_i \equiv 7 (\mathrm{mod} 8), \\ p \equiv 3 (\mathrm{mod} 4), \\ \Pi_i^+(D)' = 0. \end{cases}$$

Proof. The results follow easily from Proposition 2.3. $\square$

Proposition 2.5 We assume $\varepsilon = -1$ and the elliptic curve $E = E_-$ be as in (1.1).

(A) For $d \in \mathbb{Q}(S, 2)$, if one of the following conditions holds:

(1) $p \mid d$; (2) $q \mid d$; (3) $d = -1$.

Then $d \notin S^{(\varphi)}(E/\mathbb{Q})$.

(B$_1$) For $d = 2 \in \mathbb{Q}(S, 2)$, we have

(1) $C_2(\mathbb{Q}_2) \neq \emptyset \iff D(D + 2p + 2) \equiv 1 (\mathrm{mod} 16)$.

(2) For each prime number $l \mid pqD$, $C_2(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{2}{l}\right) = 1$, i.e., $l \equiv 1, 7 (\mathrm{mod} 8)$.

(B$_2$) For $d = -2 \in \mathbb{Q}(S, 2)$, we have

(1) $C_{-2}(\mathbb{Q}_2) \neq \emptyset \iff D(-D + 2p + 2) \equiv 3 (\mathrm{mod} 16)$.

(2) For each prime number $l \mid pqD$, $C_{-2}(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{-2}{l}\right) = 1$, i.e., $l \equiv 1, 3 (\mathrm{mod} 8)$.

(C$_1$) For each D_i $(1 \leq i \leq n)$, we have

(1) $C_{D_i}(\mathbb{Q}_2) \neq \emptyset \iff D_i \equiv 1 (\mathrm{mod} 4)$.

(2) For each prime number $l \mid pq\widehat{D_i}$, $C_{D_i}(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{D_i}{l}\right) = 1$.

(3) $C_{D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(\frac{-p\widehat{D_i}}{D_i}\right) = \left(\frac{-q\widehat{D_i}}{D_i}\right) = 1$.

(C$_2$) For each D_i $(1 \leq i \leq n)$, we have

(1) $C_{-D_i}(\mathbb{Q}_2) \neq \emptyset \iff D_i \equiv 3 (\mathrm{mod} 4)$.

(2) For each prime number $l \mid pq\widehat{D_i}$, $C_{-D_i}(\mathbb{Q}_l) \neq \emptyset \iff \left(\frac{-D_i}{l}\right) = 1$.

(3) $C_{-D_i}(\mathbb{Q}_{D_i}) \neq \emptyset \iff \left(\frac{p\widehat{D_i}}{D_i}\right) = \left(\frac{q\widehat{D_i}}{D_i}\right) = 1$.

Proof. The proof is similar to Proposition 2.1, we omit the details.

Corollary 2.6 We assume $\varepsilon = -1$ and the elliptic curve $E = E_-$ be as in

(1.1).

(A_1) $2 \in S^{(\varphi)}(E/\mathbb{Q}) \iff p \equiv 7(\mathrm{mod} 8)$ and $D_i \equiv 1, 7(\mathrm{mod} 8)$ $(1 \le i \le n)$.

(A_2) $-2 \in S^{(\varphi)}(E/\mathbb{Q}) \iff p \equiv 1(\mathrm{mod} 8)$ and $D_i \equiv 1, 3(\mathrm{mod} 8)$ $(1 \le i \le n)$.

(B_1) For each D_i $(1 \le i \le n)$, we have $D_i \in S^{(\varphi)}(E/\mathbb{Q})$ if and only if $D_i \equiv 1(\mathrm{mod} 4)$ and $\left(\frac{D_j}{D_i}\right) = \left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1$ $(1 \le j \le n$, and $j \ne i)$.

(B_2) For each D_i $(1 \le i \le n)$, we have $-D_i \in S^{(\varphi)}(E/\mathbb{Q})$ if and only if $D_i \equiv 3(\mathrm{mod} 4)$ and $\left(\frac{D_j}{D_i}\right) = \left(\frac{p}{D_i}\right) = \left(\frac{q}{D_i}\right) = 1$ $(1 \le j \le n$, and $j \ne i)$.

Proof. The results follow easily from Proposition 2.5. $\square$

Proposition 2.7 We assume $\varepsilon = -1$ and the elliptic curve $E' = E'_-$ be as in (1.2).

(A) (1) For any $d \in \mathbb{Q}(S, 2)$ and $d > 0$, $C'_d(\mathbb{R}) \ne \emptyset$. If $2 \mid d$ or $d < 0$, then $d \notin S^{(\widehat{\varphi})}(E'/\mathbb{Q})$.

(2) $\{1, pq, pD, qD\} \subset S^{(\widehat{\varphi})}(E'/\mathbb{Q})$.

(B) For each D_i $(1 \le i \le n)$, we have

(1) $C'_{D_i}(\mathbb{Q}_2) \ne \emptyset$ if and only if one of the following conditions holds:

(a) $D_i \equiv 1(\mathrm{mod} 8)$; (b) $(1 - p\widehat{D_i})(1 - q\widehat{D_i}) \equiv 0(\mathrm{mod} 16)$;

(c) $D_i \equiv 3(\mathrm{mod} 8)$ and $p \equiv 1(\mathrm{mod} 4)$; (d) $D_i \equiv 7(\mathrm{mod} 8)$ and $p \equiv 3(\mathrm{mod} 4)$.

(2) $C'_{D_i}(\mathbb{Q}_p) \ne \emptyset$ and $C'_{D_i}(\mathbb{Q}_q) \ne \emptyset$.

(3) For each $j \ne i$, $C'_{D_i}(\mathbb{Q}_{D_j}) \ne \emptyset \iff \left(1 - \left(\frac{D_i}{D_j}\right)\right)\left(1 - \left(\frac{pqD_i}{D_j}\right)\right) = 0$.

(4) $C'_{D_i}(\mathbb{Q}_{D_i}) \ne \emptyset \iff \left(1 - \left(\frac{p\widehat{D_i}}{D_i}\right)\right)\left(1 - \left(\frac{q\widehat{D_i}}{D_i}\right)\right) = 0$.

(C) (1) $C'_D(\mathbb{Q}_2) \ne \emptyset$ if and only if one of the following conditions holds:

(a) $D \equiv 1(\mathrm{mod} 8)$; (b) $p \equiv 1, 7(\mathrm{mod} 8)$;

(c) $D\equiv 3(\mathrm{mod} 8)$ and $p\equiv 5(\mathrm{mod} 8)$; (d) $D\equiv 7(\mathrm{mod} 8)$ and $p\equiv 3(\mathrm{mod} 8)$.

(2) $C'_D(\mathbb{Q}_p)\neq\emptyset$ and $C'_D(\mathbb{Q}_q)\neq\emptyset$.

(3) For each i, $C'_D(\mathbb{Q}_{D_i})\neq\emptyset\Longleftrightarrow \left(1-\left(\frac{p}{D_i}\right)\right)\left(1-\left(\frac{q}{D_i}\right)\right)=0$.

Proof. The proof is similar to Proposition 2.3, we omit the details.

Corollary 2.8 We assume $\varepsilon=-1$ and the elliptic curve $E'=E'_-$ be as in (1.2).

For each D_i $(1\leq i\leq n)$, denote

$$\Pi_i^-(D)'=\left(1-\left(\frac{p\widehat{D_i}}{D_i}\right)\right)\left(1-\left(\frac{q\widehat{D_i}}{D_i}\right)\right)+\sum_{j=1,\ j\neq i}^{n}\left(1-\left(\frac{D_i}{D_j}\right)\right)\left(1-\left(\frac{pqD_i}{D_j}\right)\right).$$

Then $D_i\in S^{(\widehat{\varphi})}(E'/\mathbb{Q})$ if and only if one of the following conditions holds:

$$(1)\ \begin{cases} D_i\equiv 1(\mathrm{mod} 8),\\ \Pi_i^-(D)'=0;\end{cases}\qquad (2)\ \begin{cases}(1-p\widehat{D_i})(1-q\widehat{D_i})\equiv 0(\mathrm{mod} 16),\\ \Pi_i^-(D)'=0;\end{cases}$$

$$(3)\ \begin{cases} D_i\equiv 3(\mathrm{mod} 8),\\ p\equiv 1(\mathrm{mod} 4),\\ \Pi_i^-(D)'=0;\end{cases}\qquad (4)\ \begin{cases} D_i\equiv 7(\mathrm{mod} 8),\\ p\equiv 3(\mathrm{mod} 4),\\ \Pi_i^-(D)'=0.\end{cases}$$

Proof. The results follow easily from Proposition 2.7. □

Now we come to prove the Theorems.

Proof of Theorem 1.1. (1) For each $i\in\{1,\cdots,n\}$, we have $\left[\frac{1}{1+\Pi_i^+(D)}\right]=0,\ 1$. Furthermore, $\left[\frac{1}{1+\Pi_i^+(D)}\right]=1\Longleftrightarrow\Pi_i^+(D)=0\Longleftrightarrow\left(\frac{-1}{D_i}\right)=1$ and $\left(\frac{p}{D_i}\right)=\left(\frac{q}{D_i}\right)=\left(\frac{D_j}{D_i}\right)=1$ $(1\leq j\leq n$ and $j\neq i)$ $\Longleftrightarrow D_i\in S^{(\varphi)}(E/\mathbb{Q})$ by Proposition 2.1. Then the conclusion (1) follows.

(2) Conclusion (2) follows easily from the above (1) and Corollary 2.2 (A). This proves Theorem 2.1. □

Proof of Corollary 1.2. By Proposition 2.1, we know that $d\notin S^{(\varphi)}(E/\mathbb{Q})$ for any $d\in\mathbb{Q}(S,2)$ satisfying one of the following conditions holds: (1) $d<0$;

(2) $p \mid d$; (3) $q \mid d$. Since $S^{(\varphi)}(E/\mathbb{Q}) \subset \mathbb{Q}(S,2) =< -1, 2, p, q, D_1, \cdots, D_n >$, we get $S^{(\varphi)}(E/\mathbb{Q}) \subset < 2, D_1, \cdots, D_n >$. Furthermore, if condition (A) holds, then by Proposition 2.1, we have $D_1, \cdots, D_n \in S^{(\varphi)}(E/\mathbb{Q})$, so $< D_1, \cdots, D_n > \subset S^{(\varphi)}(E/\mathbb{Q}) \subset < 2, D_1, \cdots, D_n >$. If condition (B) holds, then by Proposition 2.1, we have $D_1, \cdots, D_n \in S^{(\varphi)}(E/\mathbb{Q})$ but $2 \notin S^{(\varphi)}(E/\mathbb{Q})$, hence $S^{(\varphi)}(E/\mathbb{Q}) \cong < D_1, \cdots, D_n > \cong (\mathbb{Z}/2\mathbb{Z})^n$. If condition (C) holds, then by Proposition 2.1, we have $2, D_1, \cdots, D_n \in S^{(\varphi)}(E/\mathbb{Q})$, hence $S^{(\varphi)}(E/\mathbb{Q}) \cong < 2, D_1, \cdots, D_n > \cong (\mathbb{Z}/2\mathbb{Z})^{n+1}$. This proves Corollary 1.2. □

Proof of Theorem 1.5. The results of the Selmer groups are given in Corollaries 1.2 and 1.4 above.

Now $E(\mathbb{Q})[2] = \{O, (0,0), (-\varepsilon pD, 0), (-\varepsilon qD, 0)\}$, $\varphi(E(\mathbb{Q})[2]) = \{O, (0,0)\} = E^{'}(\mathbb{Q})[\widehat{\varphi}]$, by the exact sequences ([S] p.298, 314, 301)

$$
\begin{array}{l}
0 \longrightarrow \frac{E^{'}(\mathbb{Q})}{\varphi(E(\mathbb{Q}))} \longrightarrow S^{(\varphi)}(E/\mathbb{Q}) \longrightarrow \mathrm{TS}(E/\mathbb{Q})[\varphi] \longrightarrow 0, \\
0 \longrightarrow \frac{E(\mathbb{Q})}{\widehat{\varphi}(E'(\mathbb{Q}))} \longrightarrow S^{(\widehat{\varphi})}(E'/\mathbb{Q}) \longrightarrow \mathrm{TS}(E'/\mathbb{Q})[\widehat{\varphi}] \longrightarrow 0, \\
0 \longrightarrow \frac{E^{'}(\mathbb{Q})[\widehat{\varphi}]}{\varphi(E(\mathbb{Q})[2])} \longrightarrow \frac{E^{'}(\mathbb{Q})}{\varphi(E(\mathbb{Q}))} \longrightarrow \frac{E(\mathbb{Q})}{2E(\mathbb{Q})} \longrightarrow \frac{E(\mathbb{Q})}{\widehat{\varphi}(E^{'}(\mathbb{Q}))} \longrightarrow 0, \quad \text{we get}
\end{array}
$$

$$
\begin{aligned}
&\mathrm{rank}(E(\mathbb{Q})) + \dim_2(\mathrm{TS}(E/\mathbb{Q})[\varphi]) + \dim_2(\mathrm{TS}(E^{'}/\mathbb{Q})[\widehat{\varphi}]) \\
&= \dim_2(S^{(\varphi)}(E/\mathbb{Q})) + \dim_2(S^{(\widehat{\varphi})}(E^{'}/\mathbb{Q})) - 2.
\end{aligned}
$$

From this and Corollaries 1.2 and 1.4, the results in Theorem 1.5 follow, and the proof is completed. □

Depending on the corresponding results in Propositions 2.3, 2.5, 2.7, the Theorems 1.3, 1.6, 1.8, 1.10 and their Corollaries can be similarly proved.

References

[C] J.W.S. Cassels, Lectures on Elliptic Curves, LMS 24, Cambridge University Press, New York, 1995.

[**IK**] H. Iwaniec and E. Kowalski, Analytic Number Theory, American Mathematical Society, Providence, Rhode Island, 2004.

[**KS**] R. Kloosterman and E. F. Schaefer, Selmer groups of elliptic curves that can be arbitrarily large, Journal of Number Theory, 99: 148-163 (2003).

[**L**] S. Lang, Algebra, Revised Third Edition, GTM 211, Springer-Verlag, New York, 2002.

[**MSS**] J.R. Merriman, S. Siksek and N.P.Smart, Explicit 4-descents on an elliptic curve, Acta Arithmetica lxxvii.(4):385-404(1996).

[**QZ**] D. Qiu, X. Zhang, Mordell-weil groups and selmer groups of two types of elliptic curves, Science in China (series A), 2002, Vol.45, No.11, 1372-1380.

[**Sch**] S. Schmitt, Computation of the Selmer groups of certain parametrized elliptic curve, Acta Arithmetica lxxviii.(3):241-254(1997).

[**S**] J.H.Silverman, The Arithmetic of Elliptic Curves, GTM 106, Springer-Verlag, New York, 1986.

[**ST**] R.J.Strocker and J.Top, On the equation $y^2 = (x+p)(x^2+p^2)$, Rocky Mountain J. of Math. 24(1994), 1135-1161.